CERTIFICATE OF ANALYSIS

o-TOPOLIN-7-β-D-GLUCOPYRANOSIDE

[6-((2-hydroxybenzyl)amino)-purine-7-β-D-glucopyranoside]

$C_{18}H_{21}N_5O_6$, FW = 403.40

OLCHEMIM CODE: 001 787n
OLCHEMIM BATCH: B-787
MANUFACTURING DATE: September 2025
EXPIRATION DATE: September 2034
CAS:N/A
BATCH SIZE: 4 mg

SYNTHESIS

o-topolin-7-β-D-glucopyranoside was prepared by the condensation of 6-chloropurine-7-β-D-glucoside-tetraacetate and 2-hydroxybenzylamine. 6-chloropurine-7-β-D-glucoside-tetraacetate was prepared by alkylation method of 6-chloropurine with acetobromo-α-D-glucose followed chromatographic separation of 7-regioisomer. The crude product was purified on C18 reverse phase column 19 x 100 mm, 5 µm particles, gradient was formed from 10 mM aqueous ammonium acetate and acetonitrile, flow rate 15 mL/min. Fractions with the uniform product were lyophilized to get an off-white solid product.

APPEARANCE

An off-white, amorphous solid. CAUTION! VERY HYGROSCOPIC! Dissolves readily in aqueous alcohol at 5 mg/ml to give a clear, colourless solution.

ANALYSIS

UV (recorded from HPLC photodiode array detector): A_{max} 275 nm
HPLC-MS: (2.5 µm C18, 50 x 3.0 mm HPLC Column); MF: A, 0.01 M ammonium acetate in water / B, acetonitrile. At flow-rate 0.6 ml/min, the following binary gradient was used: 5% B 0-1 min; 5-60% B 1-4 min, 60-5% B 4-4.5 min. Product gives a single, symmetrical peak, Rf 2.32 min. Purity 96+%.
HRMS (HESI *m/z*): $[M+H]^+$ calcd for $C_{18}H_{22}N_5O_6$ 404.1565; found 404.1567.

STORAGE & STABILITY

ACID-LABILE! HYGROSCOPIC! Store over silica gel in tightly sealed container at - 20°C.

SAFETY

No toxicological data available for this compound. Treat as toxic and potentially mutagenic.

FOR LABORATORY USE BY EXPERIENCED PERSONNEL ONLY; NOT FOR DRUG, HOUSEHOLD OR OTHER USES

OlChemIm s.r.o., Šlechtitelů 27, P.O.Box 22, 770 10 Olomouc, Czech Republic
Tel./Fax: +420 585 427 407
email:info@olchemim.cz, OlChemIm on the Internet: http://www.olchemim.cz